



### ACTIN FILAMENTS




**Actin filaments** (also known as *microfilaments*) are two-stranded helical polymers of the protein actin. They appear as flexible structures, with a diameter of 5–9 nm, and they are organized into a variety of linear bundles, two-dimensional networks, and three-dimensional gels. Although actin filaments are dispersed throughout the cell, they are most highly concentrated in the *cortex*, just beneath the plasma membrane.

Micrographs courtesy of Roger Craig (i and iv); P.T. Matsudaira and D.R. Burgess (ii); Keith Burridge (iii).




### MICROTUBULES




**Microtubules** are long, hollow cylinders made of the protein tubulin. With an outer diameter of 25 nm, they are much more rigid than actin filaments. Microtubules are long and straight and typically have one end attached to a single microtubule-organizing center (MTOC) called a *centrosome*.

Micrographs courtesy of Richard Wade (i); D.T. Woodrow and R.W. Linck (ii); David Shima (iii); A. Desai (iv).



### INTERMEDIATE FILAMENTS



**Intermediate filaments** are ropelike fibers with a diameter of around 10 nm; they are made of intermediate filament proteins, which constitute a large and heterogeneous family. One type of intermediate filament forms a meshwork called the nuclear lamina just beneath the inner nuclear membrane. Other types extend across the cytoplasm, giving cells mechanical strength. In an epithelial tissue, they span the cytoplasm from one cell-cell junction to another, thereby strengthening the entire epithelium.

Micrographs courtesy of Roy Quinlan (i); Nancy L. Kedersha (ii); Mary Osborn (iii); Ueli Aebi (iv).

